

Anti-KRT16 antibody

Cat. No.	ml260179
Package	25 μl/100 μl/200 μl
Storage	-20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol
Product overview	
Description	Anti-KRT16 rabbit polyclonal antibody
Applications	ELISA, WB, IHC
Immunogen	Synthetic peptide of human KRT16
Reactivity	Human, Mouse, Rat
Content	0.6 mg/ml
Host species	Rabbit
lg class	Immunogen-specific rabbit IgG
Purification	Antigen affinity purification
	Antigen affinity purification
Target information	oroc
Symbol	KRT16
Full name	keratin 16
	COOL



Synonyms K16; CK16; K1CP; NEPPK; FNEPPK; KRT16A

Swissprot P08779

Target Background

The protein encoded by this gene is a member of the keratin gene family. The keratins are intermediate filament proteins responsible for the structural integrity of epithelial cells and are subdivided into cytokeratins and hair keratins. Most of the type I cytokeratins consist of acidic proteins which are arranged in pairs of heterotypic keratin chains and are clustered in a region of chromosome 17q12-q21. This keratin has been coexpressed with keratin 14 in a number of epithelial tissues, including esophagus, tongue, and hair follicles. Mutations in this gene are associated with type 1 pachyonychia congenita, non-epidermolytic palmoplantar keratoderma and unilateral palmoplantar verrucous nevus.



订购热线: 4008-898-798


Applications


Immunohistochemistry

Predicted cell location: Cytoplasm

Positive control: Human colon cancer

Recommended dilution: 50-200







Good elisa Nit producers

The image on the left is immunohistochemistry of paraffin-embedded Human colon cancer tissue using ml260179(KRT16 Antibody) at dilution 1/30, on the right is treated with synthetic peptide. (Original magnification: ×200) Predicted cell location: Cytoplasm Positive control: Human liver cancer

Recommended dilution: 50-200





The image on the left is immunohistochemistry of paraffin-embedded Human liver cancer tissue using ml260179(KRT16 Antibody) at dilution 1/30, on the right is treated with synthetic peptide. (Original magnification: ×200)

Western blotting

Predicted band size:51 kDa

Positive control:Hela cells

Recommended dilution: 500-2000



Gel: 8%SDS-PAGE Lysate: 40 µg Lane: Hela cells Primary antibody: ml260179(KRT16 Antibody) at dilution 1/450 Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution Exposure time: 20 seconds

ELISA

Recommended dilution: 2000-5000

联系电话: 4008-898-798,021-61725725

联系QQ: 2881505695, 2881505696

邮箱: mlbio_cn@yeah.net

网址: www.mlbio.cn