

# Anti-DDX56 antibody

**Cat. No.** ml263683

**Package** 25 μl/100 μl/200 μl

**Storage** -20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol

**Product overview** 

**Description** Anti-DDX56 rabbit polyclonal antibody

Applications ELISA, WB, IHC

Immunogen Synthetic peptide of human DDX56

Reactivity Human

Content 0.3 mg/ml

Host species Rabbit

Ig class Immunogen-specific rabbit IgG

Purification Antigen affinity purification

Target information

Symbol DDX56

Full name DEAD-box helicase 56



Synonyms DDX21; DDX26; NOH61

Swissprot Q9NY93

### **Target Background**

This gene encodes a member of the DEAD box protein family. DEAD box proteins, characterized by the conserved motif Asp-Glu-Ala-Asp (DEAD), are putative RNA helicases. They are implicated in a number of cellular processes involving alteration of RNA secondary structure such as translation initiation, nuclear and mitochondrial splicing, and ribosome and spliceosome assembly. Based on their distribution patterns, some members of this family are believed to be involved in embryogenesis, spermatogenesis, and cellular growth and division. The protein encoded by this gene shows ATPase activity in the presence of polynucleotides and associates with nucleoplasmic 65S preribosomal particles. This gene may be involved in ribosome synthesis, most likely during assembly of the large 60S ribosomal subunit. Multiple transcript variants encoding different isoforms have been found for this gene.




## **Applications**


# **Immunohistochemistry**

Predicted cell location: Nucleus

Positive control: Human lung cancer

Recommended dilution: 20-100





Good elisakit producere

The image on the left is immunohistochemistry of paraffin-embedded Human lung cancer tissue using ml263683(DDX56 Antibody) at dilution 1/25, on the right is treated with synthetic peptide. (Original magnification: ×200)

### Western blotting

Predicted band size:62 kDa

Positive control:HEPG2, 293T and Hela cell lysates

Recommended dilution: 200-1000



Gel: 8%SDS-PAGE


Lysate: 40 µg

Lane 1-3: HEPG2, 293T and Hela cell lysates

Primary antibody: ml263683(DDX56 Antibody) at dilution 1/400

Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution

Exposure time: 7 seconds



## **ELISA**

Recommended dilution: 5000-10000

联系电话: 4008-898-798, 021-61725725

联系QQ: 2881505695, 2881505696

邮箱: mlbio\_cn@yeah.net

网址: www.mlbio.cn