

Anti-HIST1H2BA antibody

Cat. No.	ml263694
Package	25 µl/100 µl/200 µl
Storage	-20°C, pH7.4 PBS, 0.05% NaN ₃ , 40% Glycerol

Product overview

Description	Anti-HIST1H2BA rabbit polyclonal antibody
Applications	ELISA, WB, IHC
Immunogen	Synthetic peptide of human HIST1H2BA
Reactivity	Human, Mouse, Rat
Content	1.98 mg/ml
Host species	Rabbit
Ig class	Immunogen-specific rabbit IgG
Purification	Antigen affinity purification

Target information

Symbol	HIST1H2BA
Full name	histone cluster 1 H2B family member a

Synonyms STBP; TH2B; H2BFU; TSH2B; TSH2B.1; bA317E16.3

Swissprot Q96A08

Target Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a testis/sperm-specific member of the histone H2B family. Transcripts from this gene contain a palindromic termination element.

Applications

Immunohistochemistry

Predicted cell location: Nucleus





Predicted cell location: Nucleus

Positive control: Human colorectal cancer

Positive control: Human tonsil

Recommended dilution: 50-300

Recommended dilution: 50-300



The image on the left is immunohistochemistry of paraffin-embedded Human colorectal cancer tissue using m1263694(HIST1H2BA Antibody) at dilution 1/45, on the right is treated with synthetic peptide. (Original magnification: $\times 200$)

The image on the left is immunohistochemistry of paraffin-embedded Human tonsil tissue using m1263694(HIST1H2BA Antibody) at dilution 1/45, on the right is treated with synthetic peptide. (Original magnification: $\times 200$)

Western blotting

Predicted band size: 14 kDa

Positive control: Human prostate tissue, PC-3, A549 and TM4 cell lysates

Recommended dilution: 500-2000

订购热线: 4008-898-798

Gel: 12%SDS-PAGE


Lysate: 40 μ g

Lane 1-4: Human prostate tissue, PC-3, A549 and TM4 cell lysates

Primary antibody: ml263694(HIST1H2BA Antibody) at dilution 1/500

Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution

Exposure time: 3 seconds



ELISA

Recommended dilution: 5000-10000

联系电话: 4008-898-798, 021-61725725

联系QQ: 2881505695, 2881505696

邮箱: mlbio_cn@yeah.net

网址: www.mlbio.cn