

订购热线: 4008-898-798

Anti-FBXO9 antibody

Cat. No.	ml263891
Package	25 µl/100 µl/200 µl
Storage	-20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol
Product overview	
Description	Anti-FBXO9 rabbit polyclonal antibody
Applications	ELISA, WB, IHC
Immunogen	Synthetic peptide of human FBXO9
Reactivity	Human, Mouse, Rat
Content	0.54 mg/ml
Host species	Rabbit
lg class	Immunogen-specific rabbit IgG
Purification	Antigen affinity purification
	AUCON
Target information	Antigen affinity purification
Symbol	FBX09
Full name	F-box protein 9
C	
	J



Synonyms FBX9

FBX9; VCIA1; NY-REN-57; dJ341E18.2

Swissprot Q9UK97

Target Background

This gene encodes a member of the F-box protein family which is characterized by an approximately 40 amino acid motif, the F-box. The F-box proteins constitute one of the four subunits of the ubiquitin protein ligase complex called SCFs (SKP1cullin-F-box), which function in phosphorylation-dependent ubiquitination. The F-box proteins are divided into 3 classes: Fbws containing WD-40 domains, Fbls containing leucine-rich repeats, and Fbxs containing either different protein-protein interaction modules or no recognizable motifs. The protein encoded by this gene belongs to the Fbxs class. Alternative splicing of this gene generates at least 3 transcript variants diverging at the 5' terminus.



订购热线: 4008-898-798


Applications

Immunohistochemistry

Predicted cell location: Cytoplasm

Positive control: Human tonsil

Recommended dilution: 10-50



The image on the left is immunohistochemistry of paraffin-embedded Human tonsil tissue using ml263891(FBXO9 Antibody) at dilution 1/20, on the right is treated with synthetic peptide. (Original magnification: ×200)

Good elisakit producers

Western blotting

Predicted band size:52 kDa

Positive control:Human fetal liver tissue lysate

Recommended dilution: 500-2000



Gel: 8%SDS-PAGE		
	kDa 130 —	
Lysate: 40 µg	95	
	72 —	
Lane: Human fetal liver tissue lysate	55 —	
Primary antibody: ml263891(FBXO9 Antibody) at dilution 1/200	36 —	
Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution	28 —	
Exposure time: 30 seconds	17 —	

ELISA

Recommended dilution: 5000-10000

联系电话: 4008-898-798,021-61725725

联系QQ: 2881505695, 2881505696

邮箱: mlbio_cn@yeah.net

网址: www.mlbio.cn