

Anti-MLST8 antibody

Cat. No.	ml263952
Package	25 µl/100 µl/200 µl
Storage	-20°C, pH7.4 PBS, 0.05% NaN ₃ , 40% Glycerol

Product overview

Description	Anti-MLST8 rabbit polyclonal antibody
Applications	ELISA, WB
Immunogen	Synthetic peptide of human MLST8
Reactivity	Human, Mouse, Rat
Content	1.44 mg/ml
Host species	Rabbit
Ig class	Immunogen-specific rabbit IgG
Purification	Antigen affinity purification

Target information

Symbol	MLST8
Full name	MTOR associated protein, LST8 homolog
Synonyms	GBL; LST8; POP3; WAT1; GbetaL
Swissprot	Q9BVC4

Target Background

Subunit of both mTORC1 and mTORC2, which regulates cell growth and survival in response to nutrient and hormonal signals. mTORC1 is activated in response to growth factors or amino acids. Growth factor-stimulated mTORC1 activation involves a AKT1-mediated phosphorylation of TSC1-TSC2, which leads to the activation of the RHEB GTPase that potently activates the protein kinase activity of mTORC1. Amino acid-signaling to mTORC1 requires its relocalization to the lysosomes mediated by the Ragulator complex and the Rag GTPases. Activated mTORC1 up-regulates protein synthesis by phosphorylating key regulators of mRNA translation and ribosome synthesis.

订购热线: 4008-898-798

Applications

Western blotting

Predicted band size:36 kDa

Positive control:Hela cell lysate

Recommended dilution: 500-2000

Gel: 8%SDS-PAGE


Lysate: 40 μ g

Lane: Hela cell lysate

Primary antibody: ml263952(MLST8 Antibody) at dilution 1/500

Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution

Exposure time: 5 seconds



ELISA

Recommended dilution:5000-10000

联系电话: 4008-898-798, 021-61725725

联系QQ: 2881505695, 2881505696

邮箱: mlbio_cn@yeah.net

网址: www.mlbio.cn