

胃脂肪酶抗体

产品货号： mIR18289

英文名称： LIPF

中文名称： 胃脂肪酶抗体

别名： Gastric lipase; Gastric triacylglycerol lipase; gL; HGL; HLAL; LIPF; LIPG_HUMAN.

研究领域： 肿瘤 细胞生物 免疫学 信号转导 新陈代谢

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Rat, Dog, Horse, Rabbit,

产品应用： ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 （石蜡切片需做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量： 45kDa

细胞定位： 分泌型蛋白

性状： Lyophilized or Liquid

浓度： 1mg/ml

免疫原： KLH conjugated synthetic peptide derived from human LIPF:101-200/398

亚 型 : IgG

纯化方法 : affinity purified by Protein A

储 存 液 : 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件 : Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

PubMed : PubMed

产品介绍 This gene encodes gastric lipase, an enzyme involved in the digestion of dietary triglycerides in the gastrointestinal tract, and responsible for 30% of fat digestion processes occurring in human. It is secreted by gastric chief cells in the fundic mucosa of the stomach, and it hydrolyzes the ester bonds of triglycerides under acidic pH conditions. The gene is a member of a conserved gene family of lipases that play distinct roles in neutral lipid metabolism. Several transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Nov 2010]

Subcellular Location:

Secreted.

Similarity:

Belongs to the AB hydrolase superfamily. Lipase family.

SWISS:

P07098

Gene ID:

8513

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.