

ADP 核糖基化因子结合蛋白 1 抗体

产品货号： mlR1477

英文名称： ARFIP1

中文名称： ADP 核糖基化因子结合蛋白 1 抗体

别名： ADP-ribosylation factor interacting protein 1; Arfaptin 1; 2310030I10Rik; Arfaptin; ARFAPTIN1; ARFIP1; AW123087; D3Ert598e; HSU52521; LOC60382; MGC117369; ADP ribosylation factor interacting protein 1; Arfaptin-1; ARFIP 1; Arfip1; ARFP1_HUMAN.

研究领域： 细胞生物 信号转导 转运蛋白

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Rat, Chicken, Dog, Pig, Cow, Horse,

产品应用： ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 IF=1:100-500 (石蜡切片需做抗原修复)
not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量： 41kDa

细胞定位： 细胞浆

性状： Lyophilized or Liquid

浓度： 1mg/ml

免疫原： KLH conjugated synthetic peptide derived from human Arfaptin 1:255-373/373

亚型： IgG

纯化方法 : affinity purified by Protein A

储存液 : 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件 : Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

PubMed : PubMed

产品介绍 : human homolog binds ADP-ribosylation factor (ARF) and inhibits ARF dependent phorbol ester induced secretion of matrix metalloproteinase 9 [RGD]

Function:

Putative target protein of ADP-ribosylation factor.

Subunit:

Interacts with non-myristoylated GTP-bound ARF3, but not to GDP-bound ARF3, and also to ARF1. Binds with lower affinity to ARF5 and with very little affinity to ARF6.

Tissue Specificity:

Ubiquitously expressed. Higher levels in liver, pancreas, placenta, skeletal muscle and heart.

Similarity:

Contains 1 AH domain.

SWISS:



P53367

Gene ID:

27236

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

交换和转运 (Trafficking and Transport)