

## 细胞分裂周期蛋白 20B 抗体

产品货号: mlR7783
英文名称: CDC20B
<b>中文名称</b> : 细胞分裂周期蛋白 20B 抗体
别名: CDC20 cell division cycle 20 homolog B; CDC20 like protein; CDC20B protein; cell division cycle homolog B (S cerevisiae); G6VTS76519; CD20B_HUMAN.
<b>研究领域:</b> 细胞生物 细胞周期蛋白 细胞分化
抗体来源: Rabbit
克隆类型 : Polyclonal
交叉反应 : Human, Mouse, Rat, Pig, Horse,
产品应用: ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 IF=1:100-500 (石蜡切片需做抗原修复)
not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.



分子量: 57kDa

细胞定位: 细胞核 细胞浆 细胞膜 细胞外基质 分泌型蛋白

性 状: Lyophilized or Liquid

浓 度: 1mg/ml

免疫原: KLH conjugated synthetic peptide derived from human CDC20B:441-519/519

亚 型: IgG

纯化方法: affinity purified by Protein A

储存液: 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件: Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20 °C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

PubMed: PubMed


产品介绍: The timely destruction of key regulators through ubiquitin-mediated proteolysis ensures the orderly progression of the cell cycle. The APC (anaphase-promoting complex) is a major component of this



degradation machinery and its activation is required for the execution of critical events. CDC20B has been implicated as a meiosis specific activator of APC and is required for sporulation in budding yeast.

Similarity:
Belongs to the WD repeat CDC20/Fizzy family.
Contains 7 WD repeats.
SWISS:
Q86Y33
Gene ID:
166979
Important Note:
This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic
applications.
产品图片





Key	Name	Parameter
	Raji-blank.043	FL1-H
_	bs-0295G-FITC(B)-Raji-2.044	FL1-H
	bs-0295P(B)-(FITC)(B#1FADA2.051	FL1-H
_	bs-7783R-(FITC)(B)-Raji-5.055	FL1-H