

丙酮酸脱氢酶激酶同功酶 3 抗体

产品货号： mIR12599

英文名称： PDK3

中文名称： 丙酮酸脱氢酶激酶同功酶 3 抗体

别 名： [Pyruvate dehydrogenase [lipoamide] kinase isozyme 3 mitochondrial; [Pyruvate dehydrogenase [lipoamide]] kinase isozyme 3; CMTX6; GS1-358P8.4; mitochondrial; PDK3; PDK3_HUMAN; Pyruvate dehydrogenase kinase isoenzyme 3; Pyruvate dehydrogenase kinase isoform 3; Pyruvate dehydrogenase kinase isozyme 3; Pyruvate dehydrogenase lipoamide kinase isozyme 3 mitochondrial.

研究领域： 细胞生物 信号转导 激酶和磷酸酶

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Rat, Dog, Pig, Horse, Rabbit, Sheep,

产品应用： ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 （石蜡切片需做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分 子 量： 47kDa

细胞定位： 细胞浆

性 状： Lyophilized or Liquid

浓 度 : 1mg/ml

免 疫 原 : KLH conjugated synthetic peptide derived from human PDK3:341-406/406

亚 型 : IgG

纯化方法 : affinity purified by Protein A

储 存 液 : 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件 : Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

PubMed : PubMed

产品介绍 : The pyruvate dehydrogenase (PDH) complex is a nuclear-encoded mitochondrial multienzyme complex that catalyzes the overall conversion of pyruvate to acetyl-CoA and CO₂. It provides the primary link between glycolysis and the tricarboxylic acid (TCA) cycle, and thus is one of the major enzymes responsible for the regulation of glucose metabolism. The enzymatic activity of PDH is regulated by a phosphorylation/dephosphorylation cycle, and phosphorylation results in inactivation of PDH. The protein encoded by this gene is one of the three pyruvate dehydrogenase kinases that inhibits the PDH complex by phosphorylation of the E1 alpha subunit. This gene is predominantly expressed in the heart and skeletal muscles. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Mar 2010]

Function:

Inhibits the mitochondrial pyruvate dehydrogenase complex by phosphorylation of the E1 alpha subunit, thus contributing to the regulation of glucose metabolism.

Subcellular Location:

Mitochondrion matrix.

Tissue Specificity:

Found almost exclusively in heart and skeletal muscle.

Similarity:

Belongs to the PDK/BCKDK protein kinase family.

Contains 1 histidine kinase domain.

SWISS:

Q15120

Gene ID:

5165

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.