

DEK 癌基因结合蛋白

产品货号： mlR0874

英文名称： DEK

中文名称： DEK 癌基因结合蛋白

别名： D6S231E; Dek; DEK gene; DEK oncogene; DEK oncogene DNA binding; DNA binding; DEK_HUMAN; OTTHUMP00000039357; Protein DEK.

研究领域： 染色质和核信号 转录调节因子 表观遗传学

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Rat, Chicken, Dog, Cow,

产品应用： WB=1:500-2000 ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 IF=1:100-500 （石蜡切片需做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量： 43kDa

细胞定位： 细胞核

性状： Lyophilized or Liquid

浓度： 1mg/ml

免疫原： KLH conjugated synthetic peptide derived from human DEK:101-200/315

亚型： IgG

纯化方法： affinity purified by Protein A

储 存 液： 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件： Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

PubMed： PubMed

产品介绍： The DEK gene encodes a protein with one SAP domain. This protein binds to cruciform and superhelical DNA which then induces positive supercoils into closed circular DNA, and is also involved during mRNA processing in splice site selection. Chromosomal aberrations involving this region, increased expression of this gene, and the presence of antibodies against this protein are all associated with various diseases. Two transcript variants encoding different isoforms have been found for this gene.

Function:

Involved in chromatin organization.

Subunit:

Found in a mRNA splicing-dependent exon junction complex (EJC) with DEK, RBM8A, RNPS1, SRRM1 and ALYREF/THOC4. Interacts with histones H2A, H2B, H3, H4, acetylated histone H4, non-phosphorylated DAXX and HDAC2. Component of the B-WICH complex, at least composed of SMARCA5/SNF2H, BAZ1B/WSTF, SF3B1, DEK, MYO1C, ERCC6, MYBBP1A and DDX21. Binds DNA.

Subcellular Location:

Nucleus. Note=Enriched in regions where chromatin is decondensed or sparse in the interphase nuclei.

Tissue Specificity:

Ubiquitous. Expressed at relatively high levels.

Post-translational modifications:

Phosphorylated by CK2. Phosphorylation fluctuates during the cell cycle with a moderate peak during G(1) phase, and weakens the binding of DEK to DNA.

DISEASE:

Note=A chromosomal aberration involving DEK is found in a subset of acute myeloid leukemia (AML); also known as acute non-lymphocytic leukemia. Translocation t(6;9)(p23;q34) with NUP214/CAN. It results in the formation of a DEK-CAN fusion gene.

Similarity:

Contains 1 SAP domain.

SWISS:

P35659

Gene ID:

7913

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.