

细胞角蛋白 6C 抗体

产品货号： mlR16828

英文名称： KRT6C

中文名称： 细胞角蛋白 6C 抗体

别名： CK-6C; CK-6E; Cytokeratin-6C; Cytokeratin-6E; K2C6C_HUMAN; K6C; K6E; keratin 6C; Keratin; Keratin K6h; Keratin type II cytoskeletal 6C; Keratin-6C; KRT6C; KRT6E; type II cytoskeletal 6C; Type-II keratin Kb12.

研究领域： 细胞生物 信号转导

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human,

产品应用： WB=1:500-2000 ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500

（石蜡切片需做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分 子 量 : 60kDa

细胞定位 : 细胞浆 细胞外基质 分泌型蛋白

性 状 : Lyophilized or Liquid

浓 度 : 1mg/ml

免 疫 原 : KLH conjugated synthetic peptide derived from human KRT6C:351-450/564

亚 型 : IgG

纯化方法 : affinity purified by Protein A

储 存 液 : 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件 : Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

PubMed : PubMed

产品介绍 background:

Keratins are intermediate filament proteins responsible for the structural integrity of epithelial cells and are subdivided into epithelial keratins and hair keratins. The type II keratins are clustered in a region of chromosome 12q13. [provided by RefSeq, Jul 2009]

Subunit:

Heterodimer of a type I and a type II keratin. KRT6 isomers associate with KRT16 and/or KRT17.

Tissue Specificity:

Constitutively expressed in distinct types of epithelia such as those in oral mucosa, esophagus, papillae of tongue and hair follicle outer root sheath.

Similarity:

Belongs to the intermediate filament family.

SWISS:

P48668

Gene ID:

286887



Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.