

14 号染色体开放阅读框 146 抗体

产品货号： mlR18356

英文名称： LRFN5/C14orf146

中文名称： 14 号染色体开放阅读框 146 抗体

别名： C14orf146; DKFZp686G0210; Fibronectin type III immunoglobulin and leucine rich repeat domains 8; FIGLER8; FLJ30803; leucine rich repeat and fibronectin type III domain containing 5; Leucine-rich repeat and fibronectin type-III domain-containing protein 5; Lrfn5; LRFN5_HUMAN; SALM5.

研究领域： 细胞生物 免疫学

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Rat, Chicken, Pig, Cow, Horse, Rabbit, Sheep,

产品应用： ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 （石蜡切片需做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量： 78kDa

细胞定位： 细胞膜

性状： Lyophilized or Liquid

浓度： 1mg/ml

免 疫 原： KLH conjugated synthetic peptide derived from human LRFN5/C14orf146:151-250/719

<Extracellular>

亚 型： IgG

纯化方法： affinity purified by Protein A

储 存 液： 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件： Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

PubMed： PubMed

产品介绍： Cell adhesion molecule that mediates homophilic cell-cell adhesion in a Ca(2+)-independent manner. Promotes neurite outgrowth in hippocampal neurons.

Subcellular Location:

Membrane.

Similarity:

Belongs to the LRFN family.

Contains 1 fibronectin type-III domain.

Contains 1 Ig-like (immunoglobulin-like) domain.

Contains 7 LRR (leucine-rich) repeats.

Contains 1 LRRCT domain.

Contains 1 LRRNT domain.

SWISS:

Q96NI6

Gene ID:

145581

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.