

FRA-1 蛋白抗体

产品货号: mlR1154

英文名称: FRA1

中文名称: FRA-1 蛋白抗体

别 名: Fos related antigen 1; FOS like antigen 1; FOS L1; FOSL 1; Fos related antigen 1; Fos-related antigen

1; FOSL1; FOSL1 protein; FOSL1_HUMAN; FRA 1; FRA-1.

研究领域: 肿瘤 细胞生物 免疫学 转录调节因子

抗体来源: Rabbit

克隆类型: Polyclonal

交叉反应 : Human, Mouse, Rat,

产品应用 : ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 (石蜡切片需

做抗原修复)

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量: 30kDa

细胞定位: 细胞核

性 状: Lyophilized or Liquid

浓 度: 1mg/ml

免疫原: KLH conjugated synthetic peptide derived from human FRA1:101-200/271

亚 型: IgG



纯化方法: affinity purified by Protein A

储存液: 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件: Store at -20 $^{\circ}$ C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20 $^{\circ}$ C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 $^{\circ}$ C.

PubMed: PubMed

产品介绍: The Fos gene family consists of 4 members: FOS, FOSB, FOSL1, and FOSL2. These genes encode leucine zipper proteins that can dimerize with proteins of the JUN family, thereby forming the transcription factor complex AP-1. As such, the FOS proteins have been implicated as regulators of cell proliferation, differentiation, and transformation. [provided by RefSeq].

Subunit:
Heterodimer.

Subcellular Location:
Nucleus.

Similarity:

Belongs to the bZIP family. Fos subfamily.

Contains 1 bZIP (basic-leucine zipper) domain.

SWISS:

P15407



ene ID:
061
nportant Note:
his product as supplied is intended for research use only not for use in human, therapoutis or diagnestic
his product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic pplications.