

B 细胞白血病淋巴瘤蛋白 7 抗体

产品货号： mlR9763

英文名称： BCL7C

中文名称： B 细胞白血病/淋巴瘤蛋白 7 抗体

别名： B cell CLL/lymphoma 7 protein family member C; B cell CLL/lymphoma 7C; B-cell CLL/lymphoma 7 protein family member C; BCL 7C; Bcl7c; BCL7C_HUMAN.

研究领域： 细胞生物 淋巴细胞 t-淋巴细胞

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Rat, Chicken, Dog, Pig, Cow,

产品应用： WB=1:500-2000 ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 IF=1:50-200 (石蜡切片需做抗原修复)

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量： 23kDa

细胞定位： 细胞核 细胞浆

性状： Lyophilized or Liquid

浓度： 1mg/ml

免疫原： KLH conjugated synthetic peptide derived from human BCL7C:9-100/217

亚型： IgG

纯化方法： affinity purified by Protein A

储存液： 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件： Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

PubMed： PubMed

产品介绍： This gene is identified by the similarity of its product to the N-terminal region of BCL7A protein. The BCL7A protein is encoded by the gene known to be directly involved in a three-way gene translocation in a Burkitt lymphoma cell line. The function of this gene has not yet been determined. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Nov 2013]

Function:

May play an anti-apoptotic role (By similarity).

Tissue Specificity:

Ubiquitous.

Similarity:

Belongs to the BCL7 family.

SWISS:

Q8WUZ0

Gene ID:

9274

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

产品图片

