

## 细胞周期蛋白依赖性激酶抑制剂 2B 抗体

产品货号： mlR20246

英文名称： CDKN2B / p15 INK4b

中文名称： 细胞周期蛋白依赖性激酶抑制剂 2B 抗体

别名： CDK inhibitory protein; CDK4B Inhibitor; CDN2B\_HUMAN; Cyclin dependent kinase 4 inhibitor B; Cyclin Dependent Kinase Inhibitor 2B; Cyclin dependent kinase inhibitor 2B p15 inhibits CDK4; Cyclin dependent kinases 4 and 6 binding protein; Cyclin-dependent kinase 4 inhibitor B; INK4B; MTS 2; MTS-2; MTS2; Multiple tumor suppressor 2; Multiple Tumor Suppressor 2; OTTHUMP00000021154; OTTHUMP00000021155; p14 CDK inhibitor; p14 INK4b; p14-INK4b; P15; p15 CDK inhibitor; p15 inhibits CDK4; p15-INK4b; p15INK4B; TP 15; TP15

研究领域： 肿瘤 心血管 细胞生物 免疫学 信号转导 干细胞 细胞周期蛋白 新陈代谢

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human,

产品应用： WB=1:500-2000 ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500

（石蜡切片需做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量： 15kDa

细胞定位： 细胞核

性状： Lyophilized or Liquid

浓 度 : 1mg/ml

免 疫 原 : KLH conjugated synthetic peptide derived from human CDKN2B:

亚 型 : IgG

纯化方法 : affinity purified by Protein A

储 存 液 : 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件 : Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

**PubMed :** PubMed

**产品介绍 :** p15 INK4b / MTS 2 (Mitotic Inhibitor / Suppressor Protein) is a specific inhibitor of cdk 4 / cdk 6 and is periodically expressed during the cell cycle. Reportedly, p15 INK4b expression is induced approximately 30 fold in human keratinocytes by treatment with TGF beta, suggesting that p15 INK4b may act as an effector of TGF beta mediated cell cycle arrest. p15 INK4b is mapped to chromosome 9p21, a region of frequent loss in a wide variety of cancers.

**Function:**

Interacts strongly with CDK4 and CDK6. Potent inhibitor. Potential effector of TGF-beta induced cell cycle arrest.

**Subunit:**

Heterodimer of CDKN2B with CDK4 or CDK6. Isoform 2 does not interact with CDK4 nor CDK6.

**Subcellular Location:**

Cytoplasm. Note=Also found in the nucleus.

**Tissue Specificity:**

Isoform 2 is expressed in normal (keratinocytes, fibroblasts) and tumor cell lines.

**Similarity:**

Belongs to the CDKN2 cyclin-dependent kinase inhibitor family.

Contains 4 ANK repeats.

**SWISS:**

P42772

**Gene ID:**

1030

**Important Note:**

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.