



CD99 样蛋白 2 抗体

产品货号 : mlR13742

英文名称 : CD99L2

中文名称 : CD99 样蛋白 2 抗体

别 名 : C99L2_HUMAN; CD99; CD99 antigen-like protein 2; CD99L2; MIC2-like protein 1; MIC2L1.

研究领域 : 细胞生物 淋巴细胞 t-淋巴细胞

抗体来源 : Rabbit

克隆类型 : Polyclonal

交叉反应 : Human,

产品应用 : WB=1:500-2000 ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500

(石蜡切片需做抗原修复)

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量 : 25kDa

细胞定位 : 细胞膜

性 状 : Lyophilized or Liquid

浓 度 : 1mg/ml

免 疫 原 : KLH conjugated synthetic peptide derived from human CD99L2:81-187/262 <Extracellular>

亚 型 : IgG



纯化方法 : affinity purified by Protein A

储 存 液 : 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件 : Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

PubMed : PubMed

产品介绍 : May function as a homophilic adhesion molecule. Functions in leukocyte-endothelial cell interactions during leukocyte extravasation, and in particular, at the diapedesis step. Does not seem to be involved in docking of leukocytes to the vessel wall or in lymphocyte diapedesis.

Subcellular Location:

Cell membrane. Cell junction.

Tissue Specificity:

Expressed in brain, heart, lung, liver, spleen, kidney, stomach, small intestine, skeletal muscle, ovary, thymus, testis and uterus. Lower expression seen in thymus. Expressed in E18 uterus and placenta.

Post-translational modifications:

O-glycosylated.

Similarity:

Belongs to the CD99 family.

SWISS:

Q8TCZ2

Gene ID:

83692

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

产品图片

