

卵巢癌相关基因 2 蛋白抗体

产品货号： mlR6298

英文名称： OVCA2

中文名称： 卵巢癌相关基因 2 蛋白抗体

别名： Candidate tumor suppressor in ovarian cancer 2; Candidate tumor suppressor OVCA2; ovarian cancer associated gene 2 protein; Ovarian cancer gene 2 protein; ovarian tumor suppressor candidate 2; OVCA2_HUMAN.

研究领域： 肿瘤 染色质和核信号 信号转导 转录调节因子 肿瘤细胞生物标志物

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Rat, Dog, Pig, Cow, Horse, Rabbit,

产品应用： ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 IF=1:100-500 （石蜡切片需做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量：24kDa

细胞定位：细胞核 细胞浆

性状：Lyophilized or Liquid

浓度：1mg/ml

免疫原：KLH conjugated synthetic peptide derived from human OVCA2:11-100/227

亚型：IgG

纯化方法：affinity purified by Protein A

储存液：0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件：Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

PubMed：PubMed

产品介绍 **background:**

Ubiquitously expressed protein which belongs to the UPF0483 (OVCA2) family. Encoded in an intron of the gene DPH1/OVCA1 (same strand).

Tissue Specificity:

Ubiquitously expressed.

Post-translational modifications:

Proteolytically degraded in response to RA and 4HPR treatment in a time- and dose-dependent manner in the promyelocytic leukemia cell line HL-60.

Similarity:

Belongs to the UPF0483 (OVCA2) family.

SWISS:

Q8WZ82

Gene ID:

124641

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

产品图片：

