

## ZC4H2 蛋白抗体

产品货号： mIR18532

英文名称： ZC4H2

中文名称： ZC4H2 蛋白抗体

别名： HCA127; Hepatocellular carcinoma-associated antigen 127; KIAA1166; OTTHUMP00000023432; Zc4h2; ZC4H2\_HUMAN; Zinc finger C4H2 domain-containing protein; Zinc finger, C4H2 domain containing.

研究领域： 细胞生物 免疫学 锌指蛋白 表观遗传学

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Rat, Chicken, Dog, Pig, Cow, Horse, Rabbit, Zebrafish, Sheep,

产品应用： ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 （石蜡切片需做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量： 26kDa

细胞定位： 细胞核 细胞浆

性状： Lyophilized or Liquid

浓度： 1mg/ml

**免 疫 原：** KLH conjugated synthetic peptide derived from human ZC4H2:151-224/224

**亚 型：** IgG

**纯化方法：** affinity purified by Protein A

**储 存 液：** 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

**保存条件：** Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

**PubMed：** PubMed

**产品介绍：** This gene encodes a member of the zinc finger domain-containing protein family. This family member has a C-terminal zinc finger domain that is characterized by four cysteine residues and two histidine residues, and it also includes a coiled-coil region. This protein has been detected as an autoantigen in hepatocellular carcinoma patients. This gene has been identified as a potential candidate for X-linked mental retardation. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Aug 2011]

**Similarity:**

Contains 1 C4H2-type zinc finger.

**SWISS:**

Q9NQZ6

**Gene ID:**

55906

**Important Note:**



This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.