

## 多腺苷二磷酸多聚酶 PARP9 抗体

产品货号: mlR19890

英文名称: PARP9

中文名称: 多腺苷二磷酸多聚酶 PARP9 抗体

别 名: ADP-ribosyltransferase diphtheria toxin-like 9; ARTD9; AW214463; B aggressive lymphoma protein; Bagl; BAL; BAL1; BC003281; DKFZp666B0810; DKFZp686M15238; EC 2.4.2.30; FLJ26637; FLJ35310; FLJ41418; FLJ43593; MGC:7868; MGC156739; MGC7868; OTTMUSP00000028034; PARP 9; PARP-9; Parp9; PARP9\_HUMAN; Poly (ADP ribose) polymerase family member 9; Poly [ADP-ribose] polymerase 9; RGD1307534.

研究领域: 肿瘤 细胞生物 b-淋巴细胞 表观遗传学

抗体来源: Rabbit

克隆类型: Polyclonal

交叉反应: Human, Mouse,

产品应用: ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 (石蜡切片需

做抗原修复)

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量: 97kDa

细胞定位: 细胞核 细胞浆

性 状: Lyophilized or Liquid



浓 度: 1mg/ml

免疫原: KLH conjugated synthetic peptide derived from human PARP9:601-700/854

亚型: IgG

纯化方法: affinity purified by Protein A

储存液: 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件: Store at -20  $^{\circ}$  C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20 $^{\circ}$  C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4  $^{\circ}$  C.

PubMed: PubMed