

富含半胱氨酸 CRIP2 蛋白抗体

产品货号: mlR14057

英文名称: CRIP2

中文名称: 富含半胱氨酸 CRIP2 蛋白抗体

别 名: CRIP 2; CRIP; CRIP2; CRIP2_HUMAN; CRP 2; CRP; CRP-2; CRP2; Cysteine rich intestinal protein; Cysteine rich protein 2; Cysteine-rich protein 2; ESP1; ESP1 protein; Hlp; Protein ESP1.

研究领域: 心血管 发育生物学 转运蛋白 结合蛋白 表观遗传学

抗体来源: Rabbit

克隆类型: Polyclonal

交叉反应: Human, Mouse, Rat, Horse,

产品应用: WB=1:500-2000 ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500

(石蜡切片需做抗原修复) not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量: 22kDa

细胞定位: 细胞核

性 状: Lyophilized or Liquid

浓 度: 1mg/ml

免疫原: KLH conjugated synthetic peptide derived from human CRIP2:17-120/208



Gene ID:

1397

亚 型: IgG
纯化方法 : affinity purified by Protein A
储 存 液: 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
保存条件: Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
PubMed: PubMed
产品介绍 : This gene encodes a putative transcription factor with two LIM zinc-binding domains. The encoded protein may participate in the differentiation of smooth muscle tissue. Alternative splicing results in
multiple transcript variants. [provided by RefSeq, Aug 2012]
Tissue Specificity:
Widespread tissue expression; highest levels in the heart.
Similarity:
Contains 2 LIM zinc-binding domains.
SWISS:
P52943



Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

产品图片

