

19 号染色体开放阅读框 7 抗体

产品货号： mlR18466

英文名称： ZC3H4/C19orf7

中文名称： 19 号染色体开放阅读框 7 抗体

别 名： C19orf7; KIAA1064; Zc3h4; ZC3H4_HUMAN; Zinc finger CCCH domain-containing protein 4.

研究领域： 细胞生物 转录调节因子 结合蛋白 表观遗传学

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse,

产品应用： ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 （石蜡切片需做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分 子 量： 140kDa

细胞定位： 细胞核

性 状： Lyophilized or Liquid

浓 度： 1mg/ml

免 疫 原： KLH conjugated synthetic peptide derived from human ZC3H3/C19orf7:801-900/1303

亚 型： IgG

纯化方法： affinity purified by Protein A

储 存 液： 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件： Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

PubMed： PubMed

产品介绍： This gene encodes a member of a family of CCCH (C-x8-C-x5-C-x3-H type) zinc finger domain-containing proteins. These zinc finger domains, which coordinate zinc finger binding and are characterized by three cysteine residues and one histidine residue, are nucleic acid-binding. Other family members are known to function in post-transcriptional regulation. [provided by RefSeq, Aug 2011]

Post-translational modifications:

Phosphorylated upon DNA damage, probably by ATM or ATR.

Similarity:

Contains 3 C3H1-type zinc fingers.

SWISS:

Q9UPT8

Gene ID:

23211

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.