

## 磷酸化丝裂原活化蛋白激酶 1 抗体

产品货号： mlR5414

英文名称： phospho-MEK1 + MEK2 (Ser222 + Ser226)

中文名称： 磷酸化丝裂原活化蛋白激酶 1 抗体

别名： MAP2K1(phospho S222 + S226); MAP2K1/MAP2K2(phospho S222 + S226); MEK1 + MEK2 (phospho S218 + S222 + S226); Dual specificity mitogen activated protein kinase kinase 1; ERK activator kinase 1; MAP kinase kinase 1; MAP kinase/Erk kinase 1; MAP/Erk kinase 1; Map2K1; MAPK/ERK kinase 1; MAPKK 1; MAPKK1; MEK 1; MEKK1; Mitogen activated protein kinase kinase 1; MKK 1; MKK1; PRKMK 1; PRKMK1;Protein kinase mitogen activated kinase 1 (MAP kinase kinase 1); Protein kinase mitogen activated kinase 1; Protein kinase mitogen activated, kinase 1; MP2K1\_HUMAN.

产品类型： 磷酸化抗体

研究领域： 肿瘤 免疫学 信号转导 转录调节因子 激酶和磷酸酶

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Rat, Chicken, Dog, Cow, Horse, Rabbit,

产品应用：WB=1:500-2000 ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 IF=1:100-500（石蜡切片需做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量：43kDa

细胞定位：细胞核 细胞浆

性状：Lyophilized or Liquid

浓度：1mg/ml

免疫原：KLH conjugated Synthesised phosphopeptide derived from human MEK1 around the phosphorylation site of Ser222:AN(p-S)FV

亚型：IgG

纯化方法：affinity purified by Protein A

储存液：0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件：Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted

in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**PubMed :** [PubMed](#)

#### **产品介绍 background:**

The protein encoded by this gene is a member of the dual specificity protein kinase family, which acts as a mitogen-activated protein (MAP) kinase kinase. MAP kinases, also known as extracellular signal-regulated kinases (ERKs), act as an integration point for multiple biochemical signals. This protein kinase lies upstream of MAP kinases and stimulates the enzymatic activity of MAP kinases upon wide variety of extra- and intracellular signals. As an essential component of MAP kinase signal transduction pathway, this kinase is involved in many cellular processes such as proliferation, differentiation, transcription regulation and development. [provided by RefSeq, Jul2008].

#### **Function:**

Dual specificity protein kinase which acts as an essential component of the MAP kinase signal transduction pathway. Binding of extracellular ligands such as growth factors, cytokines and hormones to their cell-surface receptors activates RAS and this initiates RAF1 activation. RAF1 then further activates the dual-specificity protein kinases MAP2K1/MEK1 and MAP2K2/MEK2. Both MAP2K1/MEK1 and MAP2K2/MEK2 function specifically in the MAPK/ERK cascade, and catalyze the concomitant phosphorylation of a threonine and a tyrosine residue in a Thr-Glu-Tyr sequence located in the extracellular signal-regulated kinases MAPK3/ERK1 and MAPK1/ERK2, leading to their activation and further transduction of the signal within the MAPK/ERK cascade. Depending on the cellular context, this pathway mediates diverse biological functions such as cell growth, adhesion, survival and differentiation, predominantly through the regulation of transcription, metabolism and cytoskeletal rearrangements. One target of the MAPK/ERK cascade is peroxisome proliferator-activated receptor gamma (PPARG), a nuclear receptor that promotes differentiation and apoptosis. MAP2K1/MEK1 has been shown to export PPARG from the nucleus. The MAPK/ERK cascade is also involved in the regulation of endosomal dynamics, including lysosome processing and endosome cycling through the perinuclear recycling compartment (PNRC), as well as in the fragmentation of the Golgi apparatus during mitosis.

#### **Subunit:**

Found in a complex with at least BRAF, HRAS1, MAP2K1, MAPK3/ERK1 and RGS14 (By similarity). Forms an heterodimer with MAP2K2/MEK2 (By similarity). Forms heterodimers with KSR2 which further dimerize to form tetramers (By similarity). Interacts with ARBB2, LAMTOR3, MAPK1/ERK2, MORG1 and RAF1 (By similarity). Interacts with PPARG and with isoform 1 of VRK2. Interacts with Yersinia yopJ. Interacts with SGK1. Interacts with BIRC6/bruce.

**Subcellular Location:**

Cytoplasm, cytoskeleton, centrosome. Cytoplasm, cytoskeleton, spindle pole body. Cytoplasm. Nucleus.  
Note=Localizes at centrosomes during prometaphase, midzone during anaphase and midbody during telophase/cytokinesis.

**Tissue Specificity:**

Widely expressed, with extremely low levels in brain.

**Post-translational modifications:**

Phosphorylation at Ser-218 and Ser-222 by MAP kinase kinase kinases (RAF or MEKK1) positively regulates kinase activity. Also phosphorylated at Thr-292 by MAPK1/ERK2 and at Ser-298 by PAK. MAPK1/ERK2 phosphorylation of Thr-292 occurs in response to cellular adhesion and leads to inhibition of Ser-298 phosphorylation by PAK.

**Similarity:**

Belongs to the protein kinase superfamily. STE Ser/Thr protein kinase family. MAP kinase kinase subfamily.

Contains 1 protein kinase domain.

**SWISS:**

P36507

**Gene ID:**

5604

**Important Note:**

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.