

剪接体相关蛋白 155 抗体

产品货号： mlR7417

英文名称： SAP155

中文名称： 剪接体相关蛋白 155 抗体

别名： Hsh 155; MDS; Pre mRNA processing 10; Pre mRNA splicing factor SF3b 155 kDa subunit; PRP 10; PRP10; PRPF 10; PRPF10; SAP 155; SAP155; SF3B 1; SF3B 1; SF3b155; Spliceosome associated protein 155; Splicing factor 3b subunit 1 155kDa; Splicing factor 3B subunit 1; SF3B1_HUMAN.

研究领域： 细胞生物 信号转导 表观遗传学

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Rat,

产品应用： WB=1:500-2000 ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500

(石蜡切片需做抗原修复)

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量： 146kDa

细胞定位： 细胞核

性状： Lyophilized or Liquid

浓度： 1mg/ml

免疫原： KLH conjugated synthetic peptide derived from human SAP155/SF3B1:1-100/1304

亚型： IgG

纯化方法 : affinity purified by Protein A

储存液 : 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件 : Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

PubMed : PubMed

产品介绍 : SAP 155 is a 1,304 amino acid member of the SF3B1 family and contains eleven HEAT repeats. Localized to nuclear speckles and also to the cytoplasm during mitosis, SAP 155 is a subunit of the SF3B splicing factor. The SF3B splicing factor is a U2 snRNP-associated protein complex essential for spliceosome assembly. SF3B contains the spliceosomal proteins SAP 49, SAP 130, SAP 145 and SAP 155. Concomitant with splicing catalysis, SAP 155 is phosphorylated at its N-terminal Thr-Pro dipeptide motifs by Dyrk1A and cyclin E/Cdk2. This modification of SAP 155 is vital for a functional spliceosome as it is an essential event in the basic splicing reaction. Due to alternative splicing events, various SAP 155 isoforms are produced.

Function:

SF3B1 is subunit 1 of the splicing factor 3b protein complex. Splicing factor 3b, together with splicing factor 3a and a 12S RNA unit, forms the U2 small nuclear ribonucleoproteins complex (U2 snRNP). The splicing factor 3b/3a complex binds pre-mRNA upstream of the intron's branch site in a sequence independent manner and may anchor the U2 snRNP to the pre-mRNA. Splicing factor 3b is also a component of the minor U12-type spliceosome. The carboxy-terminal two-thirds of subunit 1 have 22 non-identical, tandem HEAT repeats that form rod-like, helical structures.

Subunit:

Identified in the spliceosome C complex. Component of the U11/U12 snRNPs that are part of the U12-type spliceosome. Component of splicing factor SF3B which is composed of at least eight subunits; SF3B1/SAP155/SF3B155, SF3B2/SAP145/SF3B155, SF3B3/SAP130/SF3B130, SF3B4/SAP49/SF3B49, SF3B14A, PHF5A/SF3B14B, SF3B10 and SF3B125. Component of the B-WICH complex, at least composed of SMARCA5/SNF2H, BAZ1B/WSTF, SF3B1, DEK, MYO1C, ERCC6, MYBBP1A and DDX21. SF3B associates with the

splicing factor SF3A and a 12S RNA unit to form the U2 small nuclear ribonucleoproteins complex (U2 snRNP). SF3B1 interacts directly with the splicing factor U2AF. Phosphorylated form interacts with PPP1R8.

Subcellular Location:

Nucleus speckle. Note=During mitosis, transiently dispersed from the nuclear speckles to the cytoplasm.

Post-translational modifications:

Phosphorylated. Phosphorylation occurs concomitantly with the splicing catalytic steps. Phosphorylation on Thr-244, Thr-248 and Thr-313 by cyclin-dependent kinases promotes interaction with PPP1R8 during mitosis.

Similarity:

Belongs to the SF3B1 family.

Contains 11 HEAT repeats.

SWISS:

O75533

Gene ID:

23451

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

产品图片

