

ATP 结合盒转运蛋白 2 抗体

产品货号： mlR12441

英文名称： ABCF2

中文名称： ATP 结合盒转运蛋白 2 抗体

别名： 0710005O05Rik; ABC28; ABCF 2; ATP binding cassette sub family F (GCN20) member 2; DKFZp586K1823; Drr3; E430001O06; EST133090; fb04f06; fc74f03; fd58d07; HUSSY 18; MABC1; MGC128735; ABCF2_HUMAN.

研究领域： 细胞生物 信号转导 转录调节因子 转运蛋白 表观遗传学

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Rat, Pig,

产品应用： ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 (石蜡切片需做抗原修复)

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量： 71kDa

细胞定位： 细胞浆 细胞膜

性状： Lyophilized or Liquid

浓度： 1mg/ml

免疫原： KLH conjugated synthetic peptide derived from human ABCF2:301-400/623

亚型： IgG

纯化方法 : affinity purified by Protein A

储存液 : 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件 : Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

PubMed : PubMed

产品介绍 : The protein encoded by this gene is a member of the superfamily of ATP-binding cassette (ABC) transporters. ABC proteins transport various molecules across extra- and intracellular membranes. ABC genes are divided into seven distinct subfamilies (ABC1, MDR/TAP, MRP, ALD, OABP, GCN20, and White). This protein is a member of the GCN20 subfamily. Alternative splicing of this gene results in multiple transcript variants. [provided by RefSeq, Jul 2008].

Similarity:

Belongs to the ABC transporter superfamily. ABCF family. EF3 subfamily.

Contains 2 ABC transporter domains.

SWISS:

Q9UG63

Gene ID:

10061

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic

applications.

产品图片

