

锌指蛋白 420 抗体

产品货	号	: mlR7092
英文名	称	: ZNF420
中文名	称	: 锌指蛋白 420 抗体
别 / ZN420_		: APAK; ATM and p53 associated KZNF protein; FLJ32191; Zinc finger protein 420; ZNF 420
研究领	域	: 细胞生物 细胞凋亡 细胞周期蛋白 表观遗传学
抗体来	源	: Rabbit
克隆类	型	: Polyclonal
交叉反	应	: Human, Mouse, Rat, Pig, Cow, Horse, Sheep,

产品应用: ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 IF=1:100-500 (石蜡切片需做抗原修复) not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.



Good elisakit producers	格科科大型 4%

分子量: 80kDa

细胞定位: 细胞核

性 状: Lyophilized or Liquid

浓 度: 1mg/ml

免疫原: KLH conjugated synthetic peptide derived from human ZNF420:301-688/688

亚 型: IgG

纯化方法: affinity purified by Protein A

储存液: 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件: Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20 °C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

PubMed: PubMed

产品介绍 : ZNF420 contains 19 C2H2-type zinc fingers and 1 KRAB domain. It may be involved in transcriptional regulation. It is a negative regulator of p53-mediated apoptosis.



applications.

Function:
May be involved in transcriptional regulation.
Subcellular Location:
Nucleus (Potential).
Post-translational modifications:
Phosphorylated upon DNA damage, probably by ATM or ATR.
Similarity:
Contains 19 C2H2-type zinc fingers.
Contains 1 KRAB domain.
SWISS:
Q8TAQ5
Gene ID:
147923
Important Note:
This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic



产品图片

