

## X 染色体开放阅读框 32 抗体

产品货号： mlR18459R

英文名称： ZC3H12B/CXorf32

中文名称： X 染色体开放阅读框 32 抗体

别 名： Chromosome X open reading frame 32; CXorf32; MCP-induced protein 2; MCPIP2; OTTHUMP00000023434; Probable ribonuclease ZC3H12B; RP3-475B7.1; ZC12B\_HUMAN; ZC3H12B; Zinc finger CCCH domain-containing protein 12B; Zinc finger CCCH-type containing 12B.

研究领域： 细胞生物 转录调节因子 表观遗传学

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Rat, Dog, Pig, Horse, Rabbit,

产品应用： ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 （石蜡切片需做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分 子 量： 93kDa

细胞定位： 细胞核

性 状： Lyophilized or Liquid

浓 度： 1mg/ml

**免 疫 原：** KLH conjugated synthetic peptide derived from human ZC3H12B/CXorf32:121-220/836

**亚 型：** IgG

**纯化方法：** affinity purified by Protein A

**储 存 液：** 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

**保存条件：** Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

**PubMed：** PubMed

**产品介绍：** The protein encoded by this gene belongs to a family of CCCH-type zinc finger proteins that are involved in the proinflammatory activation of macrophages. The exact function of this family member is unknown, but it is thought to function as a ribonuclease. [provided by RefSeq, May 2010]

**Function:**

May function as RNase and regulate the levels of target RNA species.

**Similarity:**

Belongs to the ZC3H12 family.

Contains 1 C3H1-type zinc finger.

**SWISS:**

Q5HYM0

**Gene ID:**

340554

**Important Note:**

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.