

快速周期调节蛋白 SPDYE1 抗体

产品货号： mIR17644

英文名称： SPDYE1

中文名称： 快速周期调节蛋白 SPDYE1 抗体

别名： Putative WBSCR19-like protein 6; Ringo1; SPDE1_HUMAN; SPDYE; SPDYE1; Speedy E; Speedy homolog E1 (Xenopus laevis); Speedy protein E1; Speedy/RINGO cell cycle regulator family member E1; WBSCR19; Williams-Beuren syndrome chromosomal region 19 protein.

研究领域： 细胞生物 免疫学 细胞周期蛋白

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Rabbit,

产品应用： ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 （石蜡切片需

做抗原修复)

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分 子 量 : 41kDa

细胞定位 : 细胞核

性 状 : Lyophilized or Liquid

浓 度 : 1mg/ml

免 疫 原 : KLH conjugated synthetic peptide derived from human SPDYE1:201-300/336

亚 型 : IgG

纯化方法 : affinity purified by Protein A

储 存 液 : 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件 : Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

PubMed : PubMed

产品介绍 : SPDYE1 is located in the Williams-Beuren syndrome (WBS) critical region. WBS results from a hemizygous deletion of several genes on chromosome 7q11.23, thought to arise as a consequence of unequal crossing over between highly homologous low-copy repeat sequences flanking the deleted region.

Similarity:

Belongs to the Speedy/Ringo family.

SWISS:

Q8NFV5

Gene ID:

285955

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.