

## WD 重复蛋白 16 抗体

产品货号: mIR6750 英文名称: WDR16 中文名称: WD 重复蛋白 16 抗体 名: WD repeat containing protein 16; WD40 repeat protein up regulated in HCC; WDRPUH; WDR16\_HUMAN. 研究领域: 肿瘤 细胞生物 免疫学 信号转导 肿瘤细胞生物标志物 抗体来源: Rabbit 克隆类型: Polyclonal 交叉反应: Human, Mouse, Rat, Dog, Pig, Cow, Horse, 产品应用: WB=1:500-2000 ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 IF=1:100-500 (石蜡切片需 做抗原修复)

optimal dilutions/concentrations should be determined by the end user.

not yet tested in other applications.



产品介绍 background:

分 子	量	:	68kDa
细胞定	位	:	细胞浆
性	状	:	Lyophilized or Liquid
浓	度	:	1mg/ml
免 疫	原	:	KLH conjugated synthetic peptide derived from human WDR16:401-500/620
NÉ	型	:	IgG
纯化方	法	:	affinity purified by Protein A
储 存	液	:	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
	n te	mp	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable the erature for at least one month and for greater than a year when kept at -20 °C. When reconstitute 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
PubMe	ed :	Р	ubMed



WD repeat containing proteins, such as WDR16, play crucial roles in a wide range of physiologic functions, including signal transduction, RNA processing, remodeling the cytoskeleton, regulation of vesicular traffic, and cell division.

Function:
May play an essential role in the growth or survival of hepatocellular carcinoma (HCC).
iviay piay an essential role in the growth of survival of nepatocential carcinoma (nec.).
Subcellular Location:
Cytoplasm.
Tissue Specificity:
Highly expressed in testis. Up-regulated in hepatocellular carcinoma (HCC).
Similarity:
Contains 11 WD repeats.
SWISS:
Q8N1V2
Gene ID:
146845
Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic

applications.



## 产品图片:

