

## 配体依赖核受体共抑制因子样蛋白抗体

产品货号: mlR18199

英文名称: LCORL

中文名称: 配体依赖核受体共抑制因子样蛋白抗体

别 名: LCoR-like protein; LCORL; LCORL\_HUMAN; ligand dependent nuclear receptor corepressor like; Ligand-dependent nuclear receptor corepressor-like protein; MBLK1 related protein; MLR1; Transcription factor MLR1; Transcription factor MLR1.

研究领域: 细胞生物 转录调节因子 表观遗传学

抗体来源: Rabbit

克隆类型: Polyclonal

交叉反应: Human, Mouse, Rat, Chicken, Dog, Pig, Cow, Horse, Guinea Pig,

**产品应用 :** ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 (石蜡切片需

做抗原修复)

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量: 67kDa

细胞定位: 细胞核

性 状: Lyophilized or Liquid

浓 度: 1mg/ml



免 疫  $\mathbb{R}$ : KLH conjugated synthetic peptide derived from mouse LCORL:

亚 型: IgG
纯化方法: affinity purified by Protein A
储存液: 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
保存条件: Store at -20 $^{\circ}$ C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20 $^{\circ}$ C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 $^{\circ}$ C.
PubMed: PubMed
产品介绍 : May act as transcription activator that binds DNA elements with the sequence 5'-CCCTATCGATCGATCTCTACCT-3'. May play a role in spermatogenesis.
Subcellular Location:
Nucleus.
Similarity:
Contains 1 HTH psq-type DNA-binding domain.
SWISS:
Q3U285
Gene ID:
254251



## Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.