

TRIM45 蛋白抗体

产品货号： mlR16739

英文名称： TRIM45

中文名称： TRIM45 蛋白抗体

别名： FLJ13181; RING finger protein 99; RNF99; TRI45_HUMAN; TRIM45; Tripartite motif containing 45; Tripartite motif containing protein 45; Tripartite motif-containing protein 45.

研究领域： 细胞生物 转录调节因子 锌指蛋白 表观遗传学

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Rat, Dog, Pig, Cow, Horse, Sheep,

产品应用： WB=1:500-2000 ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500
(石蜡切片需做抗原修复)

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量： 64kDa

细胞定位： 细胞核 细胞浆

性状： Lyophilized or Liquid

浓度： 1mg/ml

免疫原： KLH conjugated synthetic peptide derived from human TRIM45:101-200/580

亚 型 : IgG

纯化方法 : affinity purified by Protein A

储 存 液 : 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件 : Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

PubMed : PubMed

产品介绍 : This gene encodes a member of the tripartite motif family. The encoded protein may function as a transcriptional repressor of the mitogen-activated protein kinase pathway. Alternatively spliced transcript variants have been described.[provided by RefSeq, Mar 2009]

Function:

May act as a transcriptional repressor in mitogen-activated protein kinase signaling pathway.

Subcellular Location:

Cytoplasm. Nucleus.

Tissue Specificity:

Expressed in skeletal muscle, brain, heart and pancreas.

Similarity:

Belongs to the TRIM/RBCC family.

Contains 2 B box-type zinc fingers.

Contains 1 filamin repeat.

Contains 1 RING-type zinc finger.

SWISS:

Q9H8W5

Gene ID:

80263

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

产品图片

