

## TRIM44 蛋白抗体

产品货号： mlR16738

英文名称： TRIM44

中文名称： TRIM44 蛋白抗体

别名： DIPB; HSA249128; MC7; MGC3490; Protein DIPB; TRI44\_HUMAN; TRIM44; Tripartite motif containing 44; Tripartite motif containing protein 44; Tripartite motif-containing protein 44.

研究领域： 细胞生物 免疫学 发育生物学 表观遗传学

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Rat, Cow,

产品应用： ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 （石蜡切片需做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分 子 量 : 38kDa

细胞定位 : 细胞核

性 状 : Lyophilized or Liquid

浓 度 : 1mg/ml

免 疫 原 : KLH conjugated synthetic peptide derived from human TRIM44:201-300/344

亚 型 : IgG

纯化方法 : affinity purified by Protein A

储 存 液 : 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件 : Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

**PubMed :** PubMed

**产品介绍 background:**

This gene encodes a member of the tripartite motif (TRIM) family. The TRIM motif includes three zinc-binding domains, namely a RING, a B-box type 1 and a B-box type 2, and a coiled-coil region. [provided by RefSeq, Jul 2008]

**Function:**

May play a role in the process of differentiation and maturation of neuronal cells (By similarity). May regulate the activity of TRIM17.

**Subunit:**

Interacts (via coiled coil) with TRIM17 (via coiled coil).

**Tissue Specificity:**

Highly expressed in testis.

**Similarity:**

Contains 1 B box-type zinc finger.

**SWISS:**

Q96DX7

**Gene ID:**

54765

**Important Note:**

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.