

## TRIM26 蛋白抗体

产品货号： mlR16728

英文名称： TRIM26

中文名称： TRIM26 蛋白抗体

别名： Acid finger protein; AFP; CAM25547; TRI26\_HUMAN; RING finger protein 95; RNF95; tripartite motif containing 26; widely expressed acid zinc finger protein; Zinc finger protein 173; ZNF173.

研究领域： 细胞生物 结合蛋白 表观遗传学

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Rat, Pig, Cow, Horse, Rabbit, Sheep,

产品应用： ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 （石蜡切片需做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分 子 量 : 62kDa

细胞定位 : 细胞浆

性 状 : Lyophilized or Liquid

浓 度 : 1mg/ml

免 疫 原 : KLH conjugated synthetic peptide derived from human TRIM26:251-350/574

亚 型 : IgG

纯化方法 : affinity purified by Protein A

储 存 液 : 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件 : Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

**PubMed :** PubMed

**产品介绍 background:**

The protein encoded by this gene is a member of the tripartite motif (TRIM) family. The TRIM motif includes three zinc-binding domains, a RING, a B-box type 1 and a B-box type 2, and a coiled-coil region. The protein localizes to cytoplasmic bodies. Although the function of the protein is unknown, the RING domain suggests that the protein may have DNA-binding activity. The gene localizes to the major histocompatibility complex (MHC) class I region on chromosome 6. Alternatively spliced transcript variants encoding the same protein have been found for this gene. [provided by RefSeq, Jun 2011]

**Function:**

TRIM26 is a member of the tripartite motif (TRIM) family. The TRIM motif includes three zinc-binding domains, a RING, a B-box type 1 and a B-box type 2, and a coiled-coil region. The protein localizes to cytoplasmic bodies. Although the function of the protein is unknown, the RING domain suggests that the protein may have DNA-binding activity.

**Subcellular Location:**

cytoplasmic

**Similarity:**

Belongs to the TRIM/RBCC family.

Contains 1 B box-type zinc finger.

Contains 1 B30.2/SPRY domain.

Contains 1 RING-type zinc finger.

**SWISS:**

Q12899

**Gene ID:**

7726

**Important Note:**

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.