

## TRA2A 蛋白抗体

产品货号： mlR17179

英文名称： TRA2A

中文名称： TRA2A 蛋白抗体

别名： AWMS1; HSU53209; htra-2 alpha; Putative MAPK activating protein PM24; TRA 2 alpha; TRA 2A; TRA-2 alpha; TRA2 alpha; TRA2-alpha; Tra-2  $\alpha$ ; Tra 2  $\alpha$ ; TRA2A\_HUMAN; Tra2alpha; Transformer 2 alpha; Transformer 2 protein homolog; Transformer-2 protein homolog A; Transformer-2 protein homolog alpha.

研究领域： 细胞生物 免疫学 转录调节因子 表观遗传学

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Rat, Dog, Pig, Cow, Horse, Sheep,

产品应用： ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 （石蜡切片需

做抗原修复)

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分 子 量 : 33kDa

细胞定位 : 细胞核

性 状 : Lyophilized or Liquid

浓 度 : 1mg/ml

免 疫 原 : KLH conjugated synthetic peptide derived from human TRA2A:1-100/282

亚 型 : IgG

纯化方法 : affinity purified by Protein A

储 存 液 : 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件 : Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

**PubMed :** PubMed

**产品介绍 :** Sequence-specific RNA-binding protein which participates in the control of pre-mRNA splicing.

**Function:**

Sequence-specific RNA-binding protein which participates in the control of pre-mRNA splicing.

**Subunit:**

Binds to A3 enhancer proteins SRp75, SRp55, SRp40 and SRp30.

**Subcellular Location:**

Nucleus.

Post-translational modifications:

Phosphorylated in the RS domains.

**Similarity:**

Belongs to the splicing factor SR family.

Contains 1 RRM (RNA recognition motif) domain.

**SWISS:**

Q13595

**Gene ID:**

29896

**Important Note:**

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.