

核因子 NF κ B 激活蛋白 1 抗体

产品货号： mlR6202

英文名称： CIKS

中文名称： 核因子 NF κ B 激活蛋白 1 抗体

别名： Act 1; ACT1; Activator of cAMP responsive element modulator (CREM) in testis; Adapter protein CIKS; C6ORF4; C6ORF5; C6ORF6; Chromosome 6 Open Reading Frame 4; Chromosome 6 Open Reading Frame 5; Chromosome 6 Open Reading Frame 6; CIKS; CIKS_HUMAN; Connection to IKK and SAPK / JNK; Connection to IKK and SAPK/JNK; Nuclear Factor Kappa B Activator 1; Nuclear factor NF kappa B activator 1; Nuclear factor NF-kappa-B activator 1; TRAF3 interacting protein 2; TRAF3-interacting protein 2; TRAF3IP2.

研究领域： 染色质和核信号 信号转导 转录调节因子

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Rat, Chicken, Dog, Pig, Cow, Rabbit,

产品应用： WB=1:500-2000 ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 IF=1:100-500 （石蜡切片需做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分 子 量 : 63kDa

细胞定位 : 细胞核 细胞浆 细胞外基质 分泌型蛋白

性 状 : Lyophilized or Liquid

浓 度 : 1mg/ml

免 疫 原 : KLH conjugated synthetic peptide derived from human Nuclear Factor Kappa B Activator 1:501-574/574

亚 型 : IgG

纯化方法 : affinity purified by Protein A

储 存 液 : 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件 : Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

PubMed : PubMed

产品介绍 background:

Could be involved in the activation of both NF-kappa-B via a NF-kappa-B inhibitor kinase (IKK)-dependent mechanism and stress-activated protein kinase (SAPK)/JNK.

Subunit:

Interacts with IKBKG/NF-kappa B essential modulator, with CHUK/IKK-alpha and with IKBKB/IKK-beta. Interacts with TRAF6.

Tissue Specificity:

Widely expressed.

DISEASE:

Defects in TRAF3IP2 are the cause of susceptibility to psoriasis type 13 (PSORS13) [MIM:614070]. PSORS13 is a common, chronic inflammatory disease of the skin with multifactorial etiology. It is characterized by red, scaly plaques usually found on the scalp, elbows and knees. These lesions are caused by abnormal keratinocyte proliferation and infiltration of inflammatory cells into the dermis and epidermis.

Similarity:

Contains 1 SEFIR domain.

SWISS:

O43734

Gene ID:

10758

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

产品图片：

