

## TIGD1 蛋白抗体

产品货号： mIR19966

英文名称： TIGD1

中文名称： TIGD1 蛋白抗体

别名： EEYORE; hypothetical protein LOC200765; Jerky mouse homolog like; TIGD1; TIGD1\_HUMAN;  
Tigger transposable element derived 1; Tigger transposable element-derived protein 1.

研究领域： 细胞生物 转录调节因子 表观遗传学

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Dog,

产品应用： ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 （石蜡切片需  
做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量： 67kDa

细胞定位： 细胞核

性状： Lyophilized or Liquid

浓度： 1mg/ml

**免 疫 原：** KLH conjugated synthetic peptide derived from human TIGD1:451-550/591

**亚 型：** IgG

**纯化方法：** affinity purified by Protein A

**储 存 液：** 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

**保存条件：** Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

**PubMed：** PubMed

**产品介绍：** The protein encoded by this gene belongs to the tigger subfamily of the pogo superfamily of DNA-mediated transposons in humans. These proteins are related to DNA transposons found in fungi and nematodes, and more distantly to the Tc1 and mariner transposases. They are also very similar to the major mammalian centromere protein B. The exact function of this gene is not known. [provided by RefSeq, Jul 2008]

**Subcellular Location:**

Nucleus.

**Similarity:**

Belongs to the tigger transposable element derived protein family.

Contains 1 DDE domain.

Contains 1 HTH CENPB-type DNA-binding domain.

Contains 1 HTH psq-type DNA-binding domain.

**SWISS:**

Q96MW7

**Gene ID:**

200765

**Important Note:**

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.