

核孔蛋白 1 抗体

产品货号： mlR9625

英文名称 vNPAP1

中文名称： 核孔蛋白 1 抗体

别 名： NPAP1; NPAP 1; NPAP-1; Nuclear pore associated protein 1; C15orf2; NPAP1_HUMAN.

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human,

产品应用： WB=1:500-2000 ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500

(石蜡切片需做抗原修复)

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分 子 量： 121kDa

细胞定位： 细胞核 细胞膜

性 状： Lyophilized or Liquid

浓 度： 1mg/ml

免 疫 原： KLH conjugated synthetic peptide derived from human NPAP1:801-900/1156

亚 型： IgG

纯化方法： affinity purified by Protein A

储 存 液 : 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件 : Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

PubMed : PubMed

产品介绍 : This gene is located in the Prader-Willi syndrome region on chromosome 15. This gene is biallelically expressed in adult testis and brain but is paternally imprinted in fetal brain. Defects in this gene may be associated with Prader-Willi syndrome. [provided by RefSeq, Aug 2012]

Function:

May be involved in spermatogenesis.

Subunit:

Associates with the nuclear pore complex (NPC).

Subcellular Location:

Nucleus, nucleoplasm. Nucleus inner membrane. Note=Colocalizes with the NPC and nuclear lamins at the nuclear periphery.

Tissue Specificity:

Testis-specific in adults. In fetal brain expressed only from the paternal allele.

SWISS:

Q9NZP6

Gene ID:

23742

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

产品图片

