

## 促黄体激素诱导蛋白 5 抗体

产品货号： mlR3919

英文名称： STARD5

中文名称： 促黄体激素诱导蛋白 5 抗体

别名： StAR related lipid transfer protein 5; StARD5; START domain containing 5; START domain containing protein 5; MGC10327.

研究领域： 肿瘤 细胞生物 免疫学 生长因子和激素 转录调节因子

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Rat, Dog, Pig, Horse,

产品应用： ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 IF=1:100-500 （石蜡切片需做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量： 23kDa

细胞定位： 细胞浆

性状： Lyophilized or Liquid

浓度： 1mg/ml

免疫原： KLH conjugated synthetic peptide derived from human STARD5:51-150/213

亚型： IgG

**纯化方法：** affinity purified by Protein A

**储 存 液：** 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

**保存条件：** Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

**PubMed：** PubMed

**产品介绍：** STARD5, or START domain containing 5, belongs to a family containing a 200- to 210-amino acid STAR-related lipid transfer (START) domain. STARD5 may be involved in the intracellular transport of sterols or other lipids and may bind cholesterol or other sterols.

**Function:**

May be involved in the intracellular transport of sterols or other lipids. May bind cholesterol or other sterols.

**Similarity:**

Contains 1 START domain.

**SWISS:**

Q9NSY2

**Gene ID:**

80765

**Important Note:**



This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.