

亲环蛋白（亲环素）PPIF 抗体

产品货号： mlR7624

英文名称： Cyclophilin F

中文名称： 亲环蛋白（亲环素）PPIF 抗体

别名： Cyclophilin 3; Cyclophilin D; Cyclophilin F; Cyp D; CYP3; CypD; EC 5.2.1.8; hCyp3; Peptidyl prolyl cis trans isomerase, mitochondrial; Peptidylprolyl isomerase F (cyclophilin F); Peptidylprolyl isomerase F; PPIase; PPIF; Rotamase.

研究领域： 细胞生物 信号转导 细胞凋亡

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Rat, Chicken, Pig, Cow,

产品应用： ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 IF=1:100-500 （石蜡切片需做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量：22kDa

细胞定位：细胞浆

性状：Lyophilized or Liquid

浓度：1mg/ml

免疫原：KLH conjugated synthetic peptide derived from human Cyclophilin F:31-130/207/207

亚型：IgG

纯化方法：affinity purified by Protein A

储存液：0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件：Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

PubMed：PubMed

产品介绍：The protein encoded by this gene is a member of the peptidyl-prolyl cis-trans isomerase (PPIase) family. PPIases catalyze the cis-trans isomerization of proline imidic peptide bonds in oligopeptides and

accelerate the folding of proteins. This protein is part of the mitochondrial permeability transition pore in the inner mitochondrial membrane. Activation of this pore is thought to be involved in the induction of apoptotic and necrotic cell death.

SWISS:

P30405

Gene ID:

10105

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

产品图片：

