

SND1 蛋白抗体

产品货号: mIR12652

英文名称: SND1

中文名称: SND1 蛋白抗体

知 名: 100 kDa coactivator; EBNA 2 co activator (100kD); EBNA2 coactivator p100; p100; p100 coactivator; p100 coactivator; p100 EBNA2 co activator; SND 1; SND 1; SND1; SND1_HUMAN; Staphylococcal nuclease and tudor domain containing 1; Staphylococcal nuclease domain containing 1; Staphylococcal nuclease domain containing protein 1; Staphylococcal nuclease domain-containing protein 1; TDRD11; TRD 11; TRD11; Tudor domain containing protein 11; Tudor domain-containing protein 11.

研究领域: 细胞生物 免疫学 转录调节因子 表观遗传学

抗体来源: Rabbit

克隆类型: Polyclonal

交叉反应: Human, Mouse, Rat, Dog, Pig, Cow, Horse, Sheep, Chimpanzee,

产品应用 : ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 (石蜡切片需

做抗原修复)

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量: 102kDa

细胞定位: 细胞核 细胞浆

性 状: Lyophilized or Liquid



浓 度: 1mg/ml

免疫原: KLH conjugated synthetic peptide derived from human SND1:51-150/910

亚型: IgG

纯化方法: affinity purified by Protein A

储存液: 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件: Store at -20 $^{\circ}$ C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20 $^{\circ}$ C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 $^{\circ}$ C.

PubMed: PubMed