

## 短链脱氢酶/还原酶 SDRO 抗体

产品货号： mlR21135

英文名称： SDR-O

中文名称： 短链脱氢酶/还原酶 SDRO 抗体

别名： DR9C7\_HUMAN; FLJ16333; MGC126600; MGC126602; orphan short chain dehydrogenase / reductase; Orphan short-chain dehydrogenase/reductase; RDH-S; RDHS; retinol dehydrogenase similar protein; SDR O; SDR-O; Sdr9c7; SDRO; short chain dehydrogenase/reductase family 9C, member 7; Short-chain dehydrogenase/reductase family 9C member 7.

研究领域： 肿瘤 细胞生物 信号转导

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Rat, Pig, Horse, Rabbit,

产品应用： IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 （石蜡切片需做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分 子 量 : 35kDa

细胞定位 : 细胞浆

性 状 : Lyophilized or Liquid

浓 度 : 1mg/ml

免 疫 原 : KLH conjugated synthetic peptide derived from human SDR-O:181-280/313

亚 型 : IgG

纯化方法 : affinity purified by Protein A

储 存 液 : Preservative: 15mM Sodium Azide, Constituents: 1% BSA, 0.01M PBS, pH 7.4

保存条件 : Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

PubMed : PubMed

**产品介绍 Function:**

Displays weak conversion of all-trans-retinal to all-trans-retinol in the presence of NADH. Has apparently no steroid dehydrogenase activity.

**Subcellular Location:**

Cytoplasm. Granular distribution in the whole cell.

**Tissue Specificity:**

Highly expressed in liver.

**Similarity:**

Belongs to the short-chain dehydrogenases/reductases (SDR) family.

**SWISS:**

Q8NEX9

**Gene ID:**

121214

**Important Note:**

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

产品图片

