

宿主细胞因子 2 抗体

产品货号： mlR17358

英文名称： Host cell factor C2

中文名称： 宿主细胞因子 2 抗体

别名： C2 factor; HCF 2; HCF-2; HCF2; HCFC 2; HCFC2; HCFC2_HUMAN; Host cell factor 2.

研究领域： 细胞生物 生长因子和激素 转录调节因子 表观遗传学

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Rat, Dog, Pig, Cow, Horse, Rabbit, Sheep, Guinea Pig,

产品应用： ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 （石蜡切片需做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分 子 量 : 87kDa

细胞定位 : 细胞核 细胞浆

性 状 : Lyophilized or Liquid

浓 度 : 1mg/ml

免 疫 原 : KLH conjugated synthetic peptide derived from human Host cell factor C2:201-300/792

亚 型 : IgG

纯化方法 : affinity purified by Protein A

储 存 液 : 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件 : Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

PubMed : PubMed

产品介绍： This gene encodes one of two proteins which interact with VP16, a herpes simplex virus protein that initiates virus infection. Both the encoded protein and the original Herpes host cell factor interact with VP16 through a beta-propeller domain. The original Herpes host cell factor, however, is effective at initiating viral infection while the encoded protein is not. Transcripts of varying length due to alternative polyadenylation signals have been described. [provided by RefSeq, Jul 2008]

Subcellular Location:

Cytoplasm. Nucleus.

Tissue Specificity:

Highly expressed in testis. Detected at lower levels in spleen, thymus, prostate, ovary, small intestine and colon.

Similarity:

Contains 3 fibronectin type-III domains.

Contains 4 Kelch repeats.

SWISS:

Q9Y5Z7

Gene ID:

29915

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic



applications.