

not yet tested in other applications.

S100 钙结合蛋白 A7 抗体

产品货号:	mIR6575
英文名称:	S100A7
中文名称:	S100 钙结合蛋白 A7 抗体
别 名: Calcium bind S100A7A; S10	HID 5; Protein S100 A7; Protein S100-A7; PSOR 1; PSOR1; Psoriasin 1; Psoriasin; Psoriasin1; S100 ing protein A7; S100 calcium-binding protein A7; S100A7; S100A7c; S10A7_HUMAN; S115A_MOUSE; 00A15.
研究领域:	肿瘤 信号转导 结合蛋白 肿瘤细胞生物标志物
抗体来源:	Rabbit
克隆类型:	Polyclonal
交叉反应 :	Mouse, Rat,
产品应用: 做抗原修复》	WB=1:500-2000 ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 IF=1:100-500 (石蜡切片需



optimal dilutions/concentrations should be determined by the end user.

分子量: 11kDa

细胞定位: 细胞浆 分泌型蛋白

性 状: Lyophilized or Liquid

浓 度: 1mg/ml

免疫原: KLH conjugated synthetic peptide derived from mouse S100A7:41-108/108

亚 型: lgG

纯化方法: affinity purified by Protein A

储存液: 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件: Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20 °C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

PubMed: PubMed

产品介绍 background:



Fetal ear, skin, and tongue and human cell lines. Highly up-regulated in psoriatic epidermis. Also highly expressed in the urine of bladder squamous cell carcinoma (SCC) bearing patients.

Subunit:
Interacts with RANBP9.
Subcellular Location:
Cytoplasm. Secreted. Note=Secreted by a non-classical secretory pathway.
Tissue Specificity:
Fetal ear, skin, and tongue and human cell lines. Highly up-regulated in psoriatic epidermis. Also highly expressed
in the urine of bladder squamous cell carcinoma (SCC) bearing patients.
Similarity:
Belongs to the S-101 family.
Contains 2 EF-hand domains.
SWISS:
Q6S5I3
Gene ID:
381493
Important Note:



This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

产品图片:

