

RTDR1 蛋白抗体

产品货号： mlR18885

英文名称： RTDR1

中文名称： RTDR1 蛋白抗体

别 名： 4933431K05Rik; MCG5994, isoform CRA_b; MGC16968; Rhabdoid tumor deletion region gene 1; Rhabdoid tumor deletion region protein 1; RTDR1; RTDR1_HUMAN.

研究领域： 细胞生物 免疫学

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human,

产品应用： ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 （石蜡切片需做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分 子 量： 39kDa

细胞定位： 细胞浆

性 状： Lyophilized or Liquid

浓 度： 1mg/ml

免 疫 原： KLH conjugated synthetic peptide derived from human RTDR1:21-120/347

亚 型： IgG

纯化方法： affinity purified by Protein A

储 存 液： 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件： Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

PubMed： PubMed

产品介绍： This gene encodes a protein with no known function but with slight similarity to a yeast vacuolar protein. The gene is located in a region deleted in pediatric rhabdoid tumors of the brain, kidney and soft tissues, but mutations in this gene have not been associated with the disease. [provided by RefSeq, Jul 2008]

Tissue Specificity:

Expressed in adult cerebellum, spinal cord, spleen, skeletal muscle and some/5 out of 9 rhabdoid tumors. Detected in fetal brain, lung, liver and kidney.

SWISS:

Q9UHP6

Gene ID:

27156

Important Note:



This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.