

## 热休克转录因子家族 X 连锁 1 抗体

产品货号： mlR18448

英文名称： LW1/HSFX1

中文名称： 热休克转录因子家族 X 连锁 1 抗体

别名： hCG\_27154; heat shock transcription factor family X linked 1; Heat shock transcription factor, X-linked; HSFX1; HSFX2; LW 1; LW1.

研究领域： 细胞生物 免疫学 信号转导 转录调节因子 表观遗传学

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human,

产品应用： WB=1:500-2000 ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500  
(石蜡切片需做抗原修复)

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量： 45kDa

细胞定位： 细胞核 细胞浆

性状： Lyophilized or Liquid

浓度： 1mg/ml

免疫原： KLH conjugated synthetic peptide derived from human LW1/HSFX1:181-280/423

亚型： IgG

**纯化方法：** affinity purified by Protein A

**储 存 液：** 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

**保存条件：** Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**PubMed：** PubMed

**产品介绍：** HSFX1 (heat shock transcription factor family, X linked 1), also known as LW-1, is a 423 amino acid protein that is predominately expressed in testis. Localized to the cytoplasm, HSFX1 is thought to be involved in spermatogenesis and male fertility.

**Subcellular Location:**

Nucleus (Potential). Cytoplasm.

**Tissue Specificity:**

Testis-specific.

**Similarity:**

Belongs to the HSF family.

**SWISS:**

Q9UBD0

**Gene ID:**

100506164

**Important Note:**

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

产品图片

