

## 人 k 链恒定区抗体

产品货号： mIR4742

英文名称： Ig kappa chain C region

中文名称： 人 k 链恒定区抗体

别 名： Ig kappa chain C region; HCAK 1; HCAK1; Ig kappa chain C region; IGKC; Immunoglobulin kappa constant; Immunoglobulin kappa constant region; Immunoglobulin kappa light chain; Kappa 1 immunoglobulin light chain; kappa light chain; Km; MGC111575; MGC62011; MGC72072; MGC88770; MGC88771; MGC88809; IGKC\_HUMAN.

研究领域： 免疫学

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human,

产品应用： ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 （石蜡切片需做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分 子 量： 11kDa

细胞定位： 细胞浆 分泌型蛋白

性 状： Lyophilized or Liquid

浓 度 : 1mg/ml

免 疫 原 : KLH conjugated synthetic peptide derived from human Ig kappa chain C region:1-60/106

亚 型 : IgG

纯化方法 : affinity purified by Protein A

储 存 液 : 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件 : Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

**PubMed :** PubMed

**产品介绍 :** Immunoglobulins belong to a group of related glyco proteins which make up 20% of serum proteins. Antigens and immunoglobulins react to confer immunity to individuals. Immunoglobulins have similar structures of two identical heavy chains and two identical light chains. Both the heavy chains and the light chains are divided into constant and variable regions. The constant regions have the same amino acid sequences between all the immunoglobulin classes. The variable regions have approximately 110 amino acids with high sequence variability. The amino acid sequence of the heavy chain determines the class of an immunoglobulin. The five types of immunoglobulin heavy chains are known as: IgG, IgA, IgM, IgD, and IgE. IgG is divided into four subclasses, and IgA is divided into two subclasses. In serum IgA and IgG are monomers with a single 4 polypeptide unit; while, IgM is a pentamer. IgA may also form polymers. Kappa light chain antibody can be used for the identification of leukemias, plasmacytomas and certain non Hodgkin's lymphomas. Kappa light chain contains one immunoglobulin like domain. The EU sequence has the INV allotypic marker, Ala 45 and Val 83. The ROY sequence has the INV allotypic marker, Ala 45 and Leu 83.

**Subcellular Location:**

Cytoplasmic and Secreted.

**SWISS:**

P01834

**Gene ID:**

3514

**Important Note:**

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.