

泛酸激酶 1 抗体

产品货号： mlR8337

英文名称： PANK1

中文名称： 泛酸激酶 1 抗体

别 名： coaA; hPanK; hPanK1; PANK; PANK1; PANK1_HUMAN; PANK1a; PANK1b; PanK1beta; Pantothenate kinase 1; Pantothenate kinase; Pantothenic acid kinase 1.

研究领域： 肿瘤 心血管 细胞生物 免疫学 信号转导

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Rat, Dog, Pig, Cow, Sheep,

产品应用： WB=1:500-2000 ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 IF=1:50-200 （石蜡切片需做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量：64kDa

细胞定位：细胞浆

性状：Lyophilized or Liquid

浓度：1mg/ml

免疫原：KLH conjugated synthetic peptide derived from human PANK1:501-598/598

亚型：IgG

纯化方法：affinity purified by Protein A

储存液：0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件：Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

PubMed：PubMed

产品介绍：The pantothenate kinase (PANK) family of proteins catalyzes the first step in coenzyme A (CoA). Pantothenate kinase 1 (PANK1) is a 598 amino acid member of the pantothenate kinase family that plays a role in

the physiological regulation of the intracellular CoA concentration. Localized to the cytoplasm, PANK1 is strongly inhibited by acetyl-CoA and manyl-CoA, as well as by high concentration of non-esterified CoA (CoASH). Four known isoforms of PANK1 exist as a result of alternative splicing events. Of these isoforms, PANK1a and PANK1b have been identified as the catalytically active isoforms. Isoform PANK1a is most highly expressed in brain, heart, kidney, liver, skeletal muscle and kidney. Isoform PANK1b is detected at much lower levels in kidney, liver, brain and testis and is not detected in heart or skeletal muscle.

Function:

Plays a role in the physiological regulation of the intracellular CoA concentration (By similarity).

Subcellular Location:

Cytoplasm.

Tissue Specificity:

Isoform 1 is expressed at high levels in brain, heart, kidney, liver, skeletal muscle and testis. Isoform 2 is detected at much lower levels in kidney, liver, brain and testis and is not detected in heart or skeletal muscle.

Similarity:

Belongs to the type II pantothenate kinase family.

SWISS:

Q8TE04

Gene ID:

53354

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

产品图片

