

RNA 聚合酶 II 结合蛋白 3 抗体

产品货号： mIR18115

英文名称： RPAP3

中文名称： RNA 聚合酶 II 结合蛋白 3 抗体

别名： Hypothetical Protein FLJ21908 FLJ21908; RNA polymerase II associated protein 3; RNA polymerase II-associated protein 3; rpap3; RPAP3_HUMAN; Spag.

研究领域： 细胞生物 信号转导 转录调节因子 表观遗传学

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Rat, Rabbit,

产品应用： ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 （石蜡切片需做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量： 76kDa

细胞定位： 细胞浆

性状： Lyophilized or Liquid

浓度： 1mg/ml

免 疫 原： KLH conjugated synthetic peptide derived from human RPAP3:21-120/665

亚 型： IgG

纯化方法： affinity purified by Protein A

储 存 液： 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件： Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

PubMed： PubMed

产品介绍： This gene encodes an RNA polymerase II-associated protein. The encoded protein may function in transcriptional regulation and may also regulate apoptosis. Alternatively spliced transcript variants have been described. [provided by RefSeq, Mar 2009]

Function:

Forms an interface between the RNA polymerase II enzyme and chaperone/scaffolding protein, suggesting that it is required to connect RNA polymerase II to regulators of protein complex formation.

Similarity:

Belongs to the RPAP3 family.

Contains 7 TPR repeats.

SWISS:

Q9H6T3

Gene ID:

79657

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.