

内皮细胞凋亡 E 蛋白 CE1 抗体

产品货号： mlR18112

英文名称： HYI

中文名称： 内皮细胞凋亡 E 蛋白 CE1 抗体

别 名： Endothelial cell apoptosis protein E CE1; Endothelial cell apoptosis protein E-CE1; HT036; Hydroxypyruvate isomerase; Hydroxypyruvate isomerase homolog (E. coli); Hydroxypyruvate isomerase homolog; hyi; HYI_HUMAN; MGC20767; Putative hydroxypyruvate isomerase; RP11 506B15.5.

研究领域： 细胞生物 免疫学 信号转导 细胞凋亡

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Rat, Pig, Horse, Rabbit,

产品应用： ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 （石蜡切片需做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分 子 量： 30kDa

细胞定位： 细胞浆

性 状： Lyophilized or Liquid

浓 度： 1mg/ml

免 疫 原： KLH conjugated synthetic peptide derived from human HYI:201-277/277

亚 型： IgG

纯化方法： affinity purified by Protein A

储 存 液： 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件： Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

PubMed： PubMed

产品介绍： This gene encodes a putative hydroxypyruvate isomerase, which likely catalyzes the conversion of hydroxypyruvate to 2-hydroxy-3-oxopropanoate, and may be involved in carbohydrate transport and metabolism. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Aug 2011]

Similarity:

Belongs to the hyi family.

SWISS:

Q5T013

Gene ID:

81888

Important Note:



This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.