

艾滋病 Rev 结合蛋白 2 抗体

产品货号： mIR16816

英文名称： KRR1

中文名称： 艾滋病 Rev 结合蛋白 2 抗体

别名： HIV 1 Rev binding protein 2; HIV-1 Rev-binding protein 2; HRB2; KRR1; KRR1 small subunit processome component homolog; KRR1, small subunit (SSU) processome component, homolog (yeast); KRR1_HUMAN; Rev interacting protein 1; Rev-interacting protein 1; RIP 1; Rip-1.

研究领域： 染色质和核信号 转录调节因子 细菌及病毒 表观遗传学

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Rat, Dog, Cow, Horse, Sheep,

产品应用： ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 （石蜡切片需

做抗原修复)

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分 子 量 : 44kDa

细胞定位 : 细胞核

性 状 : Lyophilized or Liquid

浓 度 : 1mg/ml

免 疫 原 : KLH conjugated synthetic peptide derived from human KRR1:201-300/381

亚 型 : IgG

纯化方法 : affinity purified by Protein A

储 存 液 : 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件 : Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

PubMed : PubMed

产品介绍 Function:

Required for 40S ribosome biogenesis. Involved in nucleolar processing of pre-18S ribosomal RNA and ribosome assembly.

Subunit:

Component of the ribosomal small subunit (SSU) processome (By similarity). Directly interacts with HIV-1 protein VPR. Also identified in a complex with NR3C1 and HIV-1 protein VPR.

Subcellular Location:

Nucleus > nucleolus.

Similarity:

Belongs to the KRR1 family.

Contains 1 KH domain.

SWISS:

Q13601

Gene ID:

11103



Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.