

整合素 β 6 Integrin β 6 抗体

产品货号 : mIR23107

英文名称 : Integrin beta 6

中文名称 : 整合素 β 6/Integrin β 6 抗体

别 名 : Integrin beta-6; ITGB6; ITB6_HUMAN.

研究领域 : 肿瘤 免疫学 信号转导 细胞粘附分子

抗体来源 : Rabbit

克隆类型 : Polyclonal

交叉反应 : Human, Mouse, Rat, Dog, Pig, Cow, Rabbit, Sheep,

产品应用 : WB=1:500-2000 ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500
(石蜡切片需做抗原修复)

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量：114kDa

细胞定位：细胞膜

性状：Lyophilized or Liquid

浓度：1mg/ml

免疫原：KLH conjugated synthetic peptide derived from human Integrin beta 6:351-450/788 <Extracellular>

亚型：IgG

纯化方法：affinity purified by Protein A

储存液：0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件：Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

PubMed：PubMed

产品介绍：Integrins are heterodimers composed of noncovalently associated transmembrane α and β subunits. The 16 α and 8 β subunits heterodimerize to produce more than 20 different receptors. Most integrin

receptors bind ligands that are components of the extracellular matrix, including Fibronectin, collagen and vitronectin. Certain integrins can also bind to soluble ligands, such as fibrinogen, or to counterreceptors on adjacent cells such as the intracellular adhesion molecules (ICAMs), leading to aggregation of cells. Ligands serve to cross-link or cluster integrins by binding to adjacent integrin receptors; both receptor clustering and ligand occupancy are necessary for the activation of integrin-mediated responses. In addition to mediating cell adhesion and cytoskeletal organization, integrins function as signaling receptors. Signals transduced by integrins play a role in many biological processes, including cell growth, differentiation, migration and apoptosis.

Function:

Integrin alpha-V/beta-6 is a receptor for fibronectin and cytostatin. It recognizes the sequence R-G-D in its ligands. Internalisation of integrin alpha-V/beta-6 via clathrin-mediated endocytosis promotes carcinoma cell invasion.

Subunit:

Heterodimer of an alpha and a beta subunit. Beta-6 associates with alpha-V. Interacts with FLNB. Interacts with HAX1. Alpha-V/beta-6 binds to foot-and-mouth disease virus (FMDV) VP1 protein, coxsackievirus A9, coxsackievirus B1 capsid proteins and acts as a receptor for these viruses.

Subcellular Location:

Membrane; Single-pass type I membrane protein.

Similarity:

Belongs to the integrin beta chain family.

Contains 1 VWFA domain.

SWISS:

P18564

Gene ID:

3694

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

产品图片

